



TITLE:

# The Smith Problem and a Counterexample to Laitinen's Conjecture(Methods of Transformation Group Theory)

AUTHOR(S):

Morimoto, Masaharu

---

CITATION:

Morimoto, Masaharu. The Smith Problem and a Counterexample to Laitinen's Conjecture(Methods of Transformation Group Theory). 数理解析研究所講究録 2006, 1517: 25-31

ISSUE DATE:

2006-10

URL:

<http://hdl.handle.net/2433/58719>

RIGHT:

# The Smith Problem and a Counterexample to Laitinen's Conjecture

岡山大学大学院自然科学研究科 森本 雅治 (Masaharu Morimoto)

Graduate School of Natural Science and Technology

Okayama University

## 1. INTRODUCTION

In this article, we describe current study of the Smith problem. Let  $G$  be a finite group. P. A. Smith [24] asked

whether the tangential  $G$ -representations  $U$  and  $V$  at two fixed points of an arbitrary smooth  $G$ -action on a sphere with exactly two fixed points are isomorphic to each other.

So it has been referred to as the *Smith problem*. We say that  $U$  and  $V$  above are *S-equivalent*. Important breakthroughs on the problem came in

- Atiyah-Bott-Milnor [1], [8]: Affirmative in semi-free actions,
- Cappell-Shaneson [2]–[4]: Negative for  $G = C_{4m}$ ,  $m \geq 2$ , and
- T. Petrie [15]–[19]: Negative for  $G = C_{pqr} \times C_{pqr}$  where  $p, q, r$  are distinct odd primes.

Surveys of relevant study so far are given in [22], [21], [7] and [13]. In case where  $G$  is an Oliver group, E. Laitinen [7] lighted the problem again with an conjecture which would guarantee existence of nonisomorphic S-equivalent  $G$ -representations  $U$  and  $V$ . Laitinen-Pawałowski [7] and Pawałowski-Solomon [13] showed that Laitinen's conjecture is true for most Oliver groups  $G$ :

- (1) perfect groups  $G$
- (2) Oliver groups  $G$  of odd order
- (3) Oliver groups  $G$  with  $N \triangleleft G$  such that  $G/N \cong C_{pq}$ , where  $p$  and  $q$  are distinct odd primes,
- (4) nonsolvable gap groups  $G \not\cong \text{PSL}(2, 27)$ .

A finite group  $G$  is called a *gap group* if there exists a (finite dimensional) real  $G$ -representation  $V$  such that

- (1)  $\dim V^P > 2 \dim V^H$  for any subgroup  $P \subset G$  of prime power order and any subgroup  $H \subset G$  with  $P \subsetneq H$ , and
- (2)  $V^N = 0$  for any normal subgroup  $N \subset G$  such that  $|G/N|$  is a prime power.

Although various affirmative evidences have been found, it turns out that there is a counterexample [9] to Laitinen's conjecture. That is, the conjecture is invalid in the case  $G = \text{Aut}(A_6)$ . We remark that  $\text{Aut}(A_6)$  is not a 'gap group'. The counterexample indicates that the notion 'gap group' may be crucial to Laitinen's conjecture and hence relevant to the Smith problem. As for gap groups, readers can refer to [6], [11], [10], and [25].

## 2. S-, D-EQUIVALENCES AND LAITINEN'S CONJECTURE

Let  $G$  be a finite group. For a smooth  $G$ -manifold  $X$  and a  $G$ -fixed point  $x$  in  $X$ , the tangent space  $T_x(X)$  of  $X$  at  $x$  is regarded as a real  $G$ -module, namely a real  $G$ -representation space. Thus  $T_x(X)$  is referred to as the *tangential  $G$ -representation of  $X$  at  $x$* . Let  $U$  and  $V$  be (finite dimensional) real  $G$ -modules. We say that  $U$  and  $V$  are *S-equivalent*, in symbol  $U \sim_S V$ , if there exists a homotopy sphere  $\Sigma$  with smooth  $G$ -action such that  $\Sigma^G = \{a, b\}$ ,  $T_a(\Sigma) \cong U$  and  $T_b(\Sigma) \cong V$ . Such a homotopy sphere with smooth  $G$ -action is called a *2FP sphere* (two-fixed-point sphere) for  $U$  and  $V$ . It is easy to show that  $\text{res}_P^G U \cong \text{res}_P^G V$  holds if  $U \sim_S V$  and  $|P| = 2, 4$ . Sanchez further showed that  $\text{res}_P^G U \cong \text{res}_P^G V$  holds if  $U \sim_S V$  and  $P$  is a  $p$ -subgroup of  $G$  for an odd prime  $p$ .

Let  $\mathcal{P}(G)$  denote the set of all subgroups of  $G$  of prime power order, where the trivial subgroup  $\{e\}$  lies in  $\mathcal{P}(G)$ . Define  $\mathcal{P}(G)^*$  by

$$\mathcal{P}(G)^* = \mathcal{P}(G) \setminus \{P \leq G \mid 8 \text{ divides } |P|\}.$$

Let  $\text{RO}(G)$  denote the real representation ring. Define the *Smith set*  $\text{Sm}(G)$  to be

$$\{[U] - [V] \in \text{RO}(G) \mid U \sim_S V\}.$$

Let  $\text{Irr}(G)^*$  denote the set of all irreducible real  $G$ -modules of dimension  $\geq 2$ . Then  $\text{Sm}(G)$  is a subset of the submodule  $\text{RO}(G)^*$  of  $\text{RO}(G)$  generated by  $\text{Irr}(G)^*$ .

Define the homomorphism  $d_G : \text{RO}(G)^* \rightarrow \mathbb{Z}$  by

$$d_G([U] - [V]) = \dim U^G - \dim V^G,$$

and the *Petrie kernel*  $\mathcal{P}^*\text{RO}(G)^*$  to be

$$\bigcap_{P \in \mathcal{P}(G)^*} \text{Ker}[\text{res}_P^G : \text{Ker}(d_G) \rightarrow \text{RO}(P)].$$

Then we have

$$\text{Sm}(G) \subset \mathcal{P}^*\text{RO}(G)^*.$$

There exists a disk  $\Delta$  with smooth  $G$ -action such that  $\Delta^G$  consists of exactly two points if and only if  $G$  is an Oliver group. We remark that a finite group  $G$  is an Oliver group if and only if  $G$  never admits a normal series

$$P \triangleleft H \triangleleft G$$

such that  $|P|$  and  $|G/H|$  are prime powers and  $H/P$  is a cyclic group. We say that  $U$  and  $V$  are *D-equivalent*, in symbol  $U \sim_D V$ , if there exists a disk  $\Delta$  with smooth  $G$ -action such that  $\Delta^G = \{a, b\}$ ,  $T_a(\Delta) \cong U$  and  $T_b(\Delta) \cong V$ . Such a disk with smooth  $G$ -action is called an *2FP disk* for  $U$  and  $V$ . Define the *Oliver set*  $\text{Oli}(G)$  by

$$\text{Oli}(G) = \{[U] - [V] \in \text{RO}(G) \mid U \sim_D V\},$$

and the kernel  $\mathcal{P}\text{RO}(G)^*$  by

$$\mathcal{P}\text{RO}(G)^* = \bigcap_{P \in \mathcal{P}(G)} \text{Ker}[\text{res}_P^G : \text{Ker}(d_G) \rightarrow \text{RO}(P)].$$

B. Oliver [14] proved the equality

$$\text{Oli}(G) = \mathcal{P}\text{RO}(G)^*.$$

Thus the Oliver set is an additive subgroup of  $\text{RO}(G)^*$ .

For an element  $g \in G$ , let  $(g)$  denote the conjugacy class of  $g$  in the group  $G$  and let  $(g)^\pm$  denote the real conjugacy class of  $g$  in  $G$ , namely  $(g)^\pm = (g) \cup (g^{-1})$ . Let  $a_G$

denote the number of real conjugacy classes  $(g)^\pm$ ,  $g \in G$ , such that the order of  $g$  is not a power of a prime. Laitinen-Pawalowski [7] showed that if  $a_G \neq 0$  then

$$a_G - 1 = \text{rank } \mathcal{PRO}(G)^*$$

and hence

**Theorem 2.1.** *If  $\text{Oli}(G) \neq 0$  then  $a_G \geq 2$ .*

The next conjecture has been referred to as Laitinen's conjecture.

**Conjecture.** Let  $G$  be an Oliver group. If  $a_G \geq 2$  then  $\text{Sm}(G) \cap \text{Oli}(G) \neq \emptyset$ .

### 3. NEW OBSERVATIONS ON 2FP SPHERES

Recall a classical lemma.

**Lemma 3.1.** *Let  $M$  be a connected, closed smooth  $C_2$ -manifold of dimension  $\geq 1$ . If the  $C_2$ -fixed point set  $F$  of  $M$  is nonempty then  $|F| \geq 2$ .*

A simple proof of the lemma is given in [9].

Let  $G$  be a finite group. The lemma above implies the next proposition.

**Proposition 3.2.** *Let  $\Sigma$  be a 2FP sphere for  $U$  and  $V$ . If  $K$  is a subgroup of  $G$  with index 2 then  $\dim U^K = \dim V^K$ .*

For each  $[W] \in \text{Irr}(G)^*$ , we have an associated homomorphism  $f_W : \text{RO}(G)^* \rightarrow \mathbb{Z}$ ; if  $x \in \text{RO}(G)^*$  then  $f_W(x)$  is the multiplicity of  $[W]$  in  $x$ . Thus, for  $x \in \text{RO}(G)^*$ , we have

$$x = \sum_{[W] \in \text{Irr}(G)^*} f_{[W]}(x) [W].$$

Define the subset  $\text{RO}(G)_0^*$  of  $\text{RO}(G)^*$  by

$$\text{RO}(G)_0^* = \bigcap_{[W] \in \text{Irr}(G)^*, P \in \mathcal{P}(G)} ((\dim W^P) f_{[W]} \text{res}_P^G)^{-1}(0),$$

where  $(\dim W^P) f_{[W]} \text{res}_P^G$  is a map  $\text{RO}(G)^* \rightarrow \text{RO}(P)$ ;

$$(\dim W^P) f_{[W]} \text{res}_P^G(x) = (\dim W^P) f_{[W]}(x) \text{res}_P^G(x)$$

for  $x \in \text{RO}(G)^*$ . Further define the subset  $\text{SRO}(G)$  of  $\text{RO}(G)^*$  by

$$\text{SRO}(G) = \mathcal{P}^* \text{RO}(G)^* \cap \text{RO}(G)_0^*.$$

With this notation, we have the following results.

**Theorem 3.3.** *The implication  $\text{Sm}(G) \subseteq \text{SRO}(G)$  holds for any finite group  $G$ .*

**Theorem 3.4.** *If  $G = \text{Aut}(A_6)$  then  $\text{SRO}(G) = 0$  and hence  $\text{Sm}(G) = 0$ .*

Note that if  $G = \text{Aut}(A_6)$  then  $a_G = 2$  and  $\text{Sm}(G) = 0$ , which disagrees with Laitinen's conjecture.

#### 4. NEW CONJECTURES

At present, we have the following conjectures.

**Conjecture 4.1.** Let  $G$  be an Oliver gap group. If  $a_G \geq 2$  then  $\text{Sm}(G) \cap \text{Oli}(G) \neq 0$ .

**Conjecture 4.2.** Let  $G$  be an Oliver group. If  $\text{SRO}(G) \neq 0$  then  $\text{Sm}(G) \neq 0$ .

The author wishes that readers are interested in the conjectures above.

#### REFERENCES

- [1] M. F. Atiyah and R. Bott, *A Lefschetz fixed point formula for elliptic complexes: II. Applications*, Ann. of Math. **88** (1968), 451–491.
- [2] , S. E. Cappell and J. L. Shaneson, *Fixed points of periodic maps*, Proc. Nat. Acad. Sci. USA **77** (1980), 5052–5054.
- [3] S. E. Cappell and J. L. Shaneson, *Fixed points of periodic differentiable maps*, Invent. Math. **68** (1982), 1–19.
- [4] S. E. Cappell and J. L. Shaneson, *Representations at fixed points*, Group Actions on Manifolds (Boulder, Colo., 1983), pp. 151–158, Contemp. Math., 36, Amer. Math. Soc., Providence, RI, 1985.
- [5] T. Petrie and J. Randall, *Transformation Groups on Manifolds*, Marcel Dekker, Inc., New York and Basel, 1984.
- [6] E. Laitinen and M. Morimoto, *Finite groups with smooth one fixed point actions on spheres*, Forum Math. **10** (1998), 479–520.

- [7] E. Laitinen and K. Pawałowski, *Smith equivalence of representations for finite perfect groups*, Proc. Amer. Math. Soc. **127** (1999), 297–307.
- [8] J. W. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. **72** (1966), 358–426.
- [9] M. Morimoto, *Smith equivalent  $\text{Aut}(A_6)$ -representations are isomorphic*, accepted by Proc. Amer. Math. Soc., 2006.
- [10] M. Morimoto, T. Sumi and M. Yanagihara, *Finite groups possessing gap modules*, in: Geometry and Topology, Aarhus 1998, eds. K. Grove, I. H. Madsen and E. K. Pedersen, pp. 329–342, Contemp. Math. 258, Amer. Math. Soc., Providence, 2000.
- [11] M. Morimoto and M. Yanagihara *The gap condition for  $S_5$  and GAP programs*, J. Fac. Env. Sci. Tech. Okayama Univ. **1** (1996), 1–13.
- [12] R. Oliver, *Fixed point sets of groups on finite acyclic complexes*, Comment. Math. Helv. **50** (1975), 155–177.
- [13] K. Pawałowski and R. R. Solomon, *Finite Oliver groups with rank integer 0 or 1 and Smith equivalence of group modules*, Algebr. Geom. Topol. **2** (2002), 843–895 (electronic).
- [14] B. Oliver, *Fixed point sets and tangent bundles of actions on disks and euclidean spaces*, Topology **35** (1996), 583–615.
- [15] T. Petrie, *G surgery. I. A survey*, in Algebraic and geometric topology (Santa Barbara, Calif., 1977), pp. 197–233, Lecture Notes in Math., 664, Springer Verlag, Berlin-Heidelberg-New York, 1978.
- [16] T. Petrie, *Pseudoequivalences of G-manifolds*, Algebraic and Geometric Topology (Proc. Symps. Pure Math., Stanford Univ., Stanford, Calif, 1976) Part 1, pp. 169–210, Proc. Symps. Pure Math., 32, Amer. Math. Soc., Providence, RI, 1978.
- [17] T. Petrie, *Three theorems in transformation groups*, Algebraic Topology (Aarhus 1978), pp. 549–572, Lecture Notes in Math., 763, Springer Verlag, Berlin-Heidelberg-New York, 1979.
- [18] T. Petrie, *The equivariant J homomorphism and Smith equivalence of representations*, Current Trends in Algebraic Topology (London, Ont., 1981), pp. 223–233, CMS Conf. Proc. 2, Part 2, Amer. Math. Soc., Providence, RI, 1982,
- [19] T. Petrie, *Smith equivalence of representations*, Math. Proc. Cambridge Philos. Soc. **94** (1983), 61–99.
- [20] T. Petrie and J. Randall, *Transformation Groups on Manifolds*, Marcel Dekker, Inc., New York and Basel, 1984.
- [21] T. Petrie and J. Randall, *Spherical isotropy representations*, Publ. Math. IHES **62** (1985), 5–40.
- [22] T. Petrie and J. Randall, *Transformation Groups on Manifolds*, Marcel Dekker, Inc., New York and Basel, 1984.

- [23] C. U. Sanchez, *Actions of groups of odd order on compact orientable manifolds*, Proc. Amer. Math. Soc. **54** (1976), 445–448.
- [24] P. A. Smith, *New results and old problems in finite transformation groups*, Bull. Amer. Math. Soc. **66** (1960), 401–415.
- [25] T. Sumi, *Gap modules for direct product groups*, J. Math. Soc. Japan **53** (2001), 975–990.